

JP2003012686

Title:
PYRAZOLE DERIVATIVE

Abstract:

PROBLEM TO BE SOLVED: To provide a pyrazole derivative useful as an agent for the prevention and/or treatment of diabetes or its pharmacologically permissible salt.
SOLUTION: The pyrazole derivative is expressed by general formula (I) [R^{1} is H, a substituted or unsubstituted lower alkyl or a substituted or unsubstituted lower alkoxy; R^{4} is a substituted or unsubstituted lower alkyl or a substituted or unsubstituted lower alkoxy; R^{2} is gluconopyranosyl group (the hydroxy group in the gluconopyranosyl group may be protected); and R^{3} is a substituted or unsubstituted aryl or a substituted or unsubstituted aromatic heterocyclic group]. The invention further relates to a pharmacologically permissible salt of the derivative.

(19) 日本国特許庁 (J P)

(12) 公開特許公報 (A)

(11) 特許出願公開番号

特開2003-12686

(P2003-12686A)

(43) 公開日 平成15年1月15日 (2003.1.15)

(51) Int.Cl. ⁷	識別記号	F I	データベース* (参考)
C 0 7 H 17/02		C 0 7 H 17/02	4 C 0 5 7
A 6 1 K 31/7056		A 6 1 K 31/7056	4 C 0 8 6
A 6 1 P 3/10		A 6 1 P 3/10	

審査請求 未請求 請求項の数21 O L (全 16 頁)

(21) 出願番号 特願2001-200388(P2001-200388)

(22) 出願日 平成13年7月2日 (2001.7.2)

(71) 出願人 000001029

協和醗酵工業株式会社

東京都千代田区大手町1丁目6番1号

(72) 発明者 白倉 史郎

静岡県駿東郡長泉町下七狩1188 協和醗酵
工業株式会社医薬総合研究所内

(72) 発明者 伊藤 靖彦

静岡県駿東郡長泉町下七狩1188 協和醗酵
工業株式会社医薬総合研究所内

(72) 発明者 日下 浩子

静岡県駿東郡長泉町下七狩1188 協和醗酵
工業株式会社医薬総合研究所内

最終頁に続く


(54) 【発明の名称】 ピラゾール誘導体

(57) 【要約】

【課題】 本発明の目的は、糖尿病予防および／または治療剤として有用なピラゾール誘導体またはその薬理学的に許容される塩を提供することにある。

【解決手段】 一般式 (I)

【化19】



〔式中、R¹は水素原子、置換もしくは非置換の低級アルキルまたは置換もしくは非置換の低級アルコキシを表し、R⁴は置換もしくは非置換の低級アルキルまたは置換もしくは非置換の低級アルコキシを表し、R²はグルコピラノシル基（グルコピラノシル基中の水酸基は保護されていてもよい）を表し、R³は置換もしくは非置換のアリールまたは置換もしくは非置換の芳香族複素環基を表す〕で表されるピラゾール誘導体またはその薬理学的に許容される塩を提供する。

L1 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2003:36456 CAPLUS <<LOGINID::20060731>>

DN 138:90016

T1 Preparation of 3-pyrazolyl glycosides for treatment of diabetes

IN Shirakura, Shiro; Ito, Yasuhiko; Kusaka, Hiroko; Kusaka, Hideaki;
Takeshita, Kenichi; Matsumoto, Yoshiko; Abe, Masayuki; Ota, Yoshihisa;
Nomoto, Yuji

PA Kyowa Hakko Kogyo Co., Ltd., Japan

SO Jpn. Kokai Tokkyo Koho, 16 pp.


CODEN: JKXXAF

DT Patent

LA Japanese

FAN. CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 2003012686	A2	20030115	JP 2001-200388	20010702 <—
PRAI	JP 2001-200388		20010702		
OS	MARPAT 138:90016				
GI					



AB 3-Pyrazolyl glycosides, in particular 3-pyrazolyl β -D-glucopyranosides [I; R1 = H, (un)substituted lower alkyl or lower alkoxy; R4 = (un)substituted lower alkyl or lower alkoxy; R5-R8 = H, hydroxy-protecting group; when at least one of R5-R8 is a hydroxy-protecting group and R5-R8 is H and also R1 is (un)substituted lower alkyl or lower alkoxy, R3 is (un)substituted aryl or heterocyclyl; or when R5-R8 is H and R1 is H or lower alkyl, R3 is p-(un)saturated lower alkylsulfonylaryl, or substituted aryl, or (un)substituted aromatic heterocyclyl] or pharmacol. acceptable salts thereof are prepared. Also disclosed are preventives or remedies for diabetes or diabetes complications, blood sugar-lowering agents, or Na⁺-glucose cotransporter (sodium-glucose cotransporter) (SGLT) inhibitors containing the above compds. I as the active ingredients. Thus, to a solution of 4.00 g 1,2-dihydro-4-[(4-methylthiophenyl)methyl]-5-trifluoromethyl-3H-pyrazol-3-one and 14.78 g 2,3,4,6-tetra-O-acetyl- β -D-glucopyranosyl bromide in 300 mL MeCN was added 9.69 g K₂CO₃ and stirred at room temperature for 3 days to give 58% 4-[(4-methylthiophenyl)methyl]-3-[(2,3,4,6-tetra-O-acetyl- β -D-glucopyranosyl)oxy]-5-trifluoromethyl-1H-pyrazole which (908 mg) was stirred with a mixture of 15 mL ethanol and 505 aqueous K₂CO₃ at room temperature for 1 h to give 7% 4-[(4-methylthiophenyl)methyl]-3-[(β -D-glucopyranosyl)oxy]-5-trifluoromethyl-1H-pyrazole (II). To a solution of 22 mg II in 1 mL MeOH was added 7 mg m-chloroperbenzoic acid and stirred at room temperature for 4 h to give 20% 4-[(4-methylsulfinylphenyl)methyl]-3-[(β -D-glucopyranosyl)oxy]-5-trifluoromethyl-1H-pyrazole (III). In a SGLT inhibition assay, III showed IC₅₀ of 0.0466 μ M for inhibiting the uptake of [14C]AMG in proximal tubule epithelial cell lines (LLC-PK1). III at 1 mg/kg i.v. increased the urinary excretion of glucose from 502 \pm 61 μ g/2 h (control) to 62,077 \pm 10,456 μ g/2 h in male SLC SD rats.